

Awareness of General Population Regarding Diabetes Mellitus Diseases among Tabuk, KSA

¹Hyder O Mirghani, ²Nawaf H Hamdi, ³Salha A Almarhapi, ⁴Naif H Hamdi, ⁵Raid A Alshehri

¹Assistant professor of internal medicine, Endocrinology and Metabolism, Tabuk University, KSA

^{2,3,4,5} Medical students, Faculty of medicine, Tabuk University, KSA

Abstract: Diabetes mellitus is a physiological condition caused by either production of inactive insulin or inadequate production of insulin. Insulin is responsible for regulating blood sugar levels.

Objectives: This study aim to unveil how well informed the population are as far as diabetes mellitus is concerned.

Methods: A total of 400 subjects were randomly selected and interviewed using two different structured questionnaires during the period from August 30, 2016 to September 17, 2016. The questionnaire were including risk factors, symptoms and complications of type 2 diabetes mellitus as well as methods of prevention The SPSS (version 20) was used for statistical analysis.

Results: Out of 400 subjects 200 were males and 200 were female, their ages ranged from 18 to 24 years old and 16.5%(66/400) were diabetic patient. In this study, 52.5%(105/200) of the male respondents were not aware of the risk factors of diabetes mellitus. This was a larger number compared to the 40.5%(81/200) of female unaware. However, 92% of both male and female respondents taking health related courses had an idea about causes of diabetes mellitus. About 67%(268/400) of the respondents were not sure whether the condition is infectious or non-communicable. only 29.25%(117/400) of participants identified high blood pressure as major risk factor. Lifestyle and genetics were the major aspects most of the respondents were aware of causes of diabetes mellitus.

Just 9.5%(38/400) knew that some medications can cause the disease. and close to 60%(240/400) were not aware of the methods of prevention.

39.5%(158/400) knew that diabetes mellitus can result peripheral neuropathy while 30%(120/400) knew that diabetes mellitus can cause retinopathy. other complications got low response including chronic renal failure 12.5%(50/400), CHD 11%(44/400), sexual dysfunction 4.5%(18/400) and stroke 4.5%(18/400).

Conclusion: There is lack of awareness of diabetes mellitus among Saudi population. This calls for larger studies to investigate the awareness of this morbid disease among the public. Raising the awareness of the population as an important sector of the community is badly needed.

Keywords: Diabetes, Mellitus, Awareness, Population, Tabuk.

1. INTRODUCTION

Diabetes mellitus is a chronic disease with high mortality and morbidity due to its complication, macro-vascular: myocardial infarction, stroke, and peripheral arterial disease and micro-vascular like renal failure, peripheral and autonomic neuropathy. And blindness, diabetes mellitus also greatly impairs quality of life Diabetes mellitus is alarmingly increasing due to increasing risk factors.

Diabetes is due to either the pancreas not producing enough insulin or the cells of the body not responding properly to the insulin produced.[5] There are three main types of diabetes mellitus:

- Type 1 DM results from the pancreas's failure to produce enough insulin. This form was previously referred to as "insulin-dependent diabetes mellitus" (IDDM) or "juvenile diabetes". The cause is unknown.[8]

- Type 2 DM begins with insulin resistance, a condition in which cells fail to respond to insulin properly.[8] As the disease progresses a lack of insulin may also develop.[9] This form was previously referred to as "non insulin-dependent diabetes mellitus" (NIDDM) or "adult-onset diabetes". The most common cause is excessive body weight and not enough exercise.[8]
- Gestational diabetes is the third main form and occurs when pregnant women without a previous history of diabetes develop high blood-sugar levels.[8]

The Kingdom of Saudi Arabia is among countries with the highest prevalence of diabetes mellitus (23.7%)[4] .

As of 2015, an estimated 415 million people had diabetes worldwide,[11] with type 2 DM making up about 90% of the cases.[12][13] This represents 8.3% of the adult population,[13] with equal rates in both women and men.[14] As of 2014[update], trends suggested the rate would continue to rise.[15] Diabetes at least doubles a person's risk of early death.[5]

From 2012 to 2015, approximately 1.5 to 5.0 million deaths each year resulted from diabetes.[6][11]

The global economic cost of diabetes in 2014 was estimated to be US\$612 billion.[16] In the United States, diabetes cost \$245 billion in 2012.[17]

Diabetes mellitus is a preventable and controllable disease by raising the awareness of the public by its progression, it is noted that people are not putting preventive measures in spite of relatives being affected with the disease, (Scobie et al. 2009).

People with diabetes can benefit from education about the disease and treatment, good nutrition to achieve a normal body weight, and exercise, with the goal of keeping both short-term and long-term blood glucose levels within acceptable bounds. In addition, given the associated higher risks of cardiovascular disease, lifestyle modifications are recommended to control blood pressure.[10]

We are not aware of researchers who have studied the awareness of diabetes mellitus among Saudi population in Tabuk area so we conducted this research to estimate the awareness of population in Tabuk area about this spreading disease.II.

2. METHODS AND MATERIAL

This cross sectional comparative study was conducted at Tabuk city, Saudi Arabia. Four hundred students were selected randomly by choosing the 15th to come while waiting at strategic place, all health workers who agree to participate during the same period were included. Participants were invited to sign a written informed consent and then responded to a self written questionnaire. A questionnaire was used to record knowledge, attitude and preventive practices towards Diabetes Mellitus among the population in Tabuk; Subjects below 18 years old were excluded and most responses were from the North region of KSA.

3. STATISTICAL ANALYSIS USED

The ethical committee of the Faculty of Medicine approved the research, and the Statistical Package for Social Sciences (SPSS) version 20 was used for analysis and Chi-Squire was used to compare categorical data, with P-value < 0.05 considered significant.

4. RESULTS

The questions in the questionnaire were mainly based on the specific objectives of this study. Every respondent was anonymously assessed individually according to their response to specific questions. This was followed by collective analysis to come up with the clear picture of the state of awareness as far as diabetes mellitus is concerned . Out of 400 subjects 200 were males and 200 were female). 90%(360/400) of them were not working in medical field. All the subjects were over 18 years old. 63%(252/400) had a bachelor degree, 14%(56/400) had a postgraduate university degree and 23%(92/400) with high school degree. In this population 30% had chronic disease, 26.5%(106/400) had DM and 3.5%(14/400) had asthma. 33%(132/400) of tested population had previous education about Diabetes mellitus.

A. Risk Factors for diabetes mellitus:

In this study, 52.5% (105/200) of the male respondents were not aware of the risk factors of diabetes mellitus. This was a larger number compared to the 40.5% (81/200) of female unaware. However, 92% of both male and female respondents taking health related courses had an idea about causes of diabetes mellitus. About 67% (268/400) of the respondents were not sure whether the condition is infectious or non-communicable. only 29.25% (117/400) of participants identified high blood pressure as major risk factor. Lifestyle and genetics were the major aspects most of the respondents were aware of causes of diabetes mellitus. Just 9.5% (38/400) knew that some medications can cause the disease. (Table 1)

Table 1: (Diabetes mellitus awareness)

| Risk Factors | Percentage of knowledge |
|---------------------------|-------------------------|
| Age | 30.75% (123/400) |
| Genetics | 68.5% (274/400) |
| Consuming more sweets | 66% (246/400) |
| Overweight or obesity | 52.25% (209/400) |
| High blood pressure | 29.25% (117/400) |
| Lack of physical activity | 71% (284/400) |
| Certain medications | 9.5% (38/400) |

B. Methods of prevention:

57% (228/400) were aware that diabetes could be prevented. Despite being aware of the causes of diabetes mellitus, only 8% of the respondents aware of risk factors for diabetes mellitus take measures to prevent it or reduce the risk of developing the condition. 93.75% (375/400) of all respondents have no measures of preventing the condition in place. (Table 2)

As the study team learnt from the clinicians in Tabuk city hospitals, most of the persons at risk of developing the condition on the basis of genetics did not place any measures in place despite having close relatives with the condition. (Table 2)

Table 2: (preventive methods knowledge)

| Methods | percentage of knowledge |
|---------------------------------|-------------------------|
| Regular physical activity. | 72.25% (290/400) |
| Eat healthy and balanced food . | 62.5% (250/400) |
| Appropriate body weight. | 60.75% (243/400) |
| No smoking. | 48% (192/400) |
| Don't drink alcohol. | 44.75% (179/400) |

C. Signs and Symptoms:

33% (132/400) of total population know about Diabetes mellitus symptoms, 16% (64/400) received this information from social media, and 17% (68/400) from a doctor. (table 4).

Table 3 (Diabetes mellitus symptoms awareness)

| Symptom | Percentage of Knowledge |
|--|-------------------------|
| Blurry Vision. | 42.5% (170/400) |
| Increase thirst or the need to urinate. | 32.75% (131/400) |
| Feeling tired or ill. | 31.5% (126/400) |
| Recurring skin, gum or bladder infections. | 28.25% (113/400) |
| Unexpected weight loss. | 26.75% (107/400) |
| Slow healing cuts or bruises. | 24% (96/400) |
| Loss of feeling in the feet. | 44.5% (178/400) |

B. Complication of Diabetes mellitus:

39.5%(158/400) knew that diabetes mellitus can result peripheral neuropathy while 30%(120/400) knew that diabetes mellitus can cause retinopathy. And 29.5%(118/400). other complications got low response including chronic renal failure 12.5%(50/400), CHD 11%(44/400), sexual dysfunction 4.5%(18/400) and stroke 4.5%(18/400).(Table 4)

Table 4: (Diabetes mellitus complication awareness)

| Complication | Percentage of Knowledge |
|------------------------|-------------------------|
| peripheral neuropathy | 39.5%(158/400) |
| retinopathy | 30%(120/400) |
| Diabetic foot | 29.5%(118/400) |
| Chronic kidney disease | 12.5%(50/400) |
| Coronary heart disease | 11%(44/400) |
| Sexual Dysfunction | 4.5%(18/400) |
| Stroke | 4.5%(18/400) |

5. DISCUSSION

The results of this study indicate information deficiency among Tabuk population. This might be the situation in the society at large, but that would require a wider scope of the study to prove. The risk of diabetes mellitus among the study population is very high mainly due to lack of awareness. This is further increased by the fact that even those aware of the condition do not take the initiative to prevent it.

Above half of all the respondents have insufficient knowledge concerning diabetes mellitus. Lack of interest and ignorance were found to be the major obstacles to attaining information about this condition. The public health sector should take the initiative of creating awareness on the major diseases affecting the society without any measures being taken. Diabetes can be prevented and controlled after its development in individuals but only with people being aware of its progression.

6. CONCLUSION

There is lack of awareness of diabetes mellitus among general population in Tabuk city. This calls for larger studies to investigate the awareness of this morbid disease among the public. Raising the awareness of the University student as an important sector of the community is badly needed.

REFERENCES

- [1] Simpson S, Corabian P, Jacobs P, Johnson JA: The cost of major co morbidity in people with diabetes mellitus. *Can Med Assoc J* 168:1661–1667, 2003
- [2] Assmann G, Nofer JR, Schulte H: Cardiovascular risk assessment in metabolic syndrome: view from PROCAM. *Endocrinol Metab Clin North Am* 33:377–392, 2004
- [3] Vinik EJ, Hayes RP, Oglesby A, Bastyr E, Barlow B, Ford-Molvisek SL, Vinik AL. The development and validation of the Norfolk QOL-DN, a new measure of patient's perception of the effects of diabetes and diabetic neuropathy. *Diabetes Technol Ther* 2005;7:497-508
- [4] Al-Nozha MM, Al-Maatouq MA, Al-Mazrou YY, Al-Harthi SS, Arafah MR, Khalil et al. Diabetes in Saudi Arabia. *Saudi Med J* 2004 Nov;25(11):1603–10.
- [5] Shoback, edited by David G. Gardner, Dolores (2011). "Chapter 17". *Greenspan's basic & clinical endocrinology* (9th ed.). New York: McGraw-Hill Medical. ISBN 0-07-162243-8.
- [6] The top 10 causes of death Fact sheet N°310". World Health Organization. Oct 2013.
- [7] Diabetes Fact sheet N°312". WHO. October 2013. Archived from the original on 26 August 2013. Retrieved 25 March 2014.

- [8] RSSDI textbook of diabetes mellitus. (Rev. 2nd ed.). New Delhi: Jaypee Brothers Medical Publishers. 2012. p. 235. ISBN 9789350254899.
- [9] Adler AI, Stratton IM, Neil HA, Yudkin JS, Matthews DR, Cull CA, Wright AD, Turner RC, Holman RR (August 2000). "Association of systolic blood pressure with macrovascular and microvascular complications of type 2 diabetes (UKPDS 36): prospective observational study". *BMJ*. 321 (7258): 412–9. doi:10.1136/bmj.321.7258.412. PMC 27455. PMID 10938049.
- [10] "Update 2015". IDF. International Diabetes Federation. p. 13. Retrieved 21 Mar 2016.
- [11] Williams textbook of endocrinology (12th ed.). Philadelphia: Elsevier/Saunders. pp. 1371–1435. ISBN 978-1-4377-0324-5.
- [12] Shi, Yuankai; Hu, Frank B (7 June 2014). "The global implications of diabetes and cancer". *The Lancet*. 383 (9933): 1947–8. doi:10.1016/S0140-6736(14)60886-2. PMID 24910221.
- [13] Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, Shibuya K, Salomon JA, Abdalla S, Aboyans V, et al. (Dec 15, 2012). "Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010.". *Lancet*. 380 (9859): 2163–96. doi:10.1016/S0140-6736(12)61729-2. PMID 23245607.
- [14] "Annual Report 2014" (PDF). IDF. International Diabetes Federation. Retrieved 13 July 2016.
- [15] IDF DIABETES ATLAS (PDF) (6th ed.). International Diabetes Federation. 2013. p. 7. ISBN 2930229853.
- [16] American Diabetes Association (Apr 2013). "Economic costs of diabetes in the U.S. in 2012.". *Diabetes Care*. 36 (4): 1033–46. doi:10.2337/dc12-2625. PMC 3609540. PMID 23468086.